



Research



Cite this article: ur Rashid H, Khan S, Irum, Khan A, Ahmad N, Shah T, Khan K. 2024 Direct synthesis, characterization, *in vitro* and *in silico* studies of simple chalcones as potential antimicrobial and antileishmanial agents. *R. Soc. Open Sci.* **11**: 240410.
<https://doi.org/10.1098/rsos.240410>

Received: 19 March 2024

Accepted: 29 May 2024

Subject Category:

Chemistry

Subject Areas:

organic chemistry/computational chemistry/
medicinal chemistry

Keywords:

chalcone, benzaldehyde, acetophenone,
biomolecule, antimicrobial

Authors for correspondence:

Nasir Ahmad

e-mail: nasirdft@gmail.com

Khalid Khan

e-mail: drkhalidchem@yahoo.com

This article has been edited by the Royal Society of Chemistry, including the commissioning, peer review process and editorial aspects up to the point of acceptance.

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.7302563>.



Direct synthesis, characterization, *in vitro* and *in silico* studies of simple chalcones as potential antimicrobial and antileishmanial agents

Haroon ur Rashid¹, Sherwali Khan², Irum¹, Asad Khan³,
Nasir Ahmad³, Tanzeel Shah⁴ and Khalid Khan³

¹Center for Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, RS, Brazil

²Department of Chemistry, Rawalpindi Women University, 6th Road Satellite Town, Rawalpindi, Pakistan

³Department of Chemistry, Islamia College University Peshawar, Khyber Pakhtunkhwa, Pakistan

⁴Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, Khyber Pakhtunkhwa, 25120, Pakistan

HR, 0000-0003-3335-7957; KK, 0000-0002-2135-6976

Chalcone represents a vital biosynthetic scaffold owing to its numerous therapeutic effects. The present study was intended to synthesize 17 chalcone derivatives (3a–q) by direct coupling of substituted acetophenones and benzaldehyde. The target chalcones were characterized by spectroscopic analyses followed by their *in vitro* antimicrobial, and antileishmanial investigations with reference to standard drugs. The majority of the chalcones displayed good to excellent biological activities. Chalcone **3q** (1000 µg ml⁻¹) exhibited the most potent antibacterial effect with its zone of inhibition values of 30, 33 and 34 mm versus *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* respectively. The results also confirmed chalcone **3q** to be the most potent versus *Leishmania major* with the lowest IC₅₀ value of 0.59 ± 0.12 µg ml⁻¹. Chalcone **3i** (500 µg ml⁻¹) was noticed to be the most potent antifungal agent with its zone of inhibition being 29 mm against *Candida albicans*. Computational studies of chalcones **3i** and **3q** supported the preliminary *in vivo* results. The existence of the amino moiety and bromine atom on ring-A and methoxy moieties on ring-B caused better biological effects of



Research article

In silico evaluation of favipiravir-associated potential new drugs against polymerase enzyme of SARS-CoV-2

Saira^a, Khalid Khan^{a,**}, Asad Khan^a, Ateeq Khan^a, Tanzeel Shah^b, Nasir Ahmad^a, Haroon ur Rashid^c, Muhammad Zahoor^{d,*}, Riaz Ullah^e, Ahmed Bari^f, Muhammad Naveed Umar^g

^a Department of Chemistry, Islamia College Peshawar, Khyber Pakhtunkhwa, 25120, Pakistan

^b Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, Khyber Pakhtunkhwa, 25120, Pakistan

^c Center of Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, RS, Brazil

^d Department of Biochemistry, University of Malakand at Chakdara, Dir Lower, Khyber Pakhtunkhwa, Pakistan

^e Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, 11451, Saudi Arabia

^f Department of Pharmaceutical Chemistry, College of Pharmacy King Saud University, Riyadh, Saudi Arabia

^g Department of Chemistry, University of Liverpool, UK

ARTICLE INFO

Keywords:

SARS-CoV-2

RdRp

Favipiravir's derivatives

Bioinformatics

ABSTRACT

Millions of lives have been lost to the deadly SARS-CoV-2 virus. Vaccines and antiviral drugs are essential scientific tools in combating viral infections. This *in silico* study focused on the RdRp inhibitor favipiravir, exploring new analogs by substituting the fluorine atom on the pyrazine ring with both homocyclic and heterocyclic moieties. Initially, ADME and toxicity properties were assessed using SwissADME and ProTox-II online tools. Ligands L6 and L7 exhibited high bioavailability and drug-likeness compared to favipiravir. Subsequently, all new analogs were docked into the RdRp active site using AutoDock Vina, demonstrating high affinity compared to favipiravir. Based on optimal ADMET profiles and docking scores, ligands L4, L6, and L7 underwent 200 ns MDS using the CHARM 36 force field in NAMD software to validate docking results. Various trajectory analyses, including RMSD, RMSF, histograms, total number of contacts, and ligand properties, were conducted to gain insights into the interaction patterns between ligands and RdRp. All protein-ligand complexes exhibited greater stability than favipiravir throughout simulations period. This theoretical study suggests that ligands L6 and L7 could serve as lead candidates for RdRp inhibition. Cell-Based SARS-CoV-2 RdRp Activity Assay is recommended to validate these *in silico* findings.

1. Introduction

Coronaviruses induce respiratory infections in mammals such as bats, masked palm civets, and camels, as well as in avian species

* Corresponding author.

** Corresponding author.

E-mail addresses: sairaphd@icp.edu.pk (Saira), drkhalidchem@yahoo.com (K. Khan), asadkhan14ch@gmail.com (A. Khan), infoateeqjan@gmail.com (A. Khan), ferromind1989@gmail.com (T. Shah), nasirdft@gmail.com (N. Ahmad), haroongold@gmail.com (H. Rashid), mohammadzahoor@yahoo.com (M. Zahoor), rullah@ksu.edu.sa (R. Ullah), abari@ksu.edu.sa (A. Bari), m.naveed-umar@liverpool.ac.uk (M.N. Umar).

<https://doi.org/10.1016/j.heliyon.2024.e38479>

Received 2 May 2024; Received in revised form 15 July 2024; Accepted 25 September 2024

Available online 26 September 2024

2405-8440/© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



In silico evaluation of the potential anticancer effects of Viscosine and Quercetin 7-rutinoside: inhibition of DEAD-box RNA helicase DDX3X and b-cell lymphoma-extra large (Bcl-XL) activity

Soma Khan¹ · Khalid Khan¹ · Tanzeel Shah² · Haroon ur Rashid³ · Nasir Ahmad¹ · Akhtar Muhammad¹ · Muhammad Zahoor⁴ · Muhammad Naveed Umar⁵ · Riaz Ullah⁶ · Zafar Iqbal⁷

Received: 18 March 2024 / Accepted: 24 September 2024 / Published online: 8 November 2024
© The Author(s), under exclusive licence to the Institute of Chemistry, Slovak Academy of Sciences 2024

Abstract

Globally, cancer is the second leading cause of mortality, surpassed only by cardiovascular disorders. Among its various types, Non-Small Cell Lung Cancer (NSCLC) poses a significant global health risk. Chemotherapy is the most common therapeutic approach for treating cancer. However, chemotherapeutic drugs can cause serious toxicity by reducing the survival of both tumor and normal cells, thereby diminishing the survival chances of patients. Natural products extracted from medicinal plants serve as effective anticancer agents with minimal side effects. Among natural compounds, flavonoids have been reported as effective anticancer agents. In this study, two flavonoids, viscosine and quercetin 7-rutinoside, were evaluated for their antitumor potential using in silico models. These flavonoids were tested for their inhibitory activity against DEAD-box RNA helicase DDX3X (referred to as DDX3) and B-cell lymphoma-extra-large (Bcl-XL) enzymes through molecular docking and dynamic simulation analyses. The docking analysis revealed low docking scores and binding energies for both viscosine and quercetin 7-rutinoside. The overall docking results revealed that the selected flavonoids exhibited a favorable binding affinity for DDX3 and Bcl-XL. They bind firmly to the key active sites and remain stably within the pocket regions of DDX3 and Bcl-XL, thereby inhibiting their activity. This data was further supported by the RMSF and RMSD analyses. The preliminary molecular docking and simulation analyses confirm the promising inhibitory effects of viscosine and quercetin 7-rutinoside against DDX3 and Bcl-XL. Therefore, these flavonoids could serve as potential antitumor drug candidates in the future. However, further experimental studies are needed to validate their potential for clinical trials.

Keywords Flavonoids · Viscosine · Quercetin 7-rutinoside · Chemotherapy · DDX3X · Bcl-XL apoptosis

✉ Khalid Khan
drkhalidchem@yahoo.com

✉ Muhammad Zahoor
mohammadzahoorus@yahoo.com

Soma Khan
jk2018@gmail.com

Tanzeel Shah
ferromind1989@gmail.com

Haroon ur Rashid
haroongold@gmail.com

Nasir Ahmad
nasirdft@gmail.com

Akhtar Muhammad
wadikurram96@yahoo.com

Muhammad Naveed Umar
m.naveed-umar@liverpool.ac.uk

Riaz Ullah
rullah@ksu.edu.sa

Zafar Iqbal
zsyed@ksu.edu.sa

¹ Department of Chemistry, Islamia College, Peshawar 25120, Khyber Pakhtunkhwa, Pakistan

² Institute of Basic Medical Sciences, Khyber Medical University, Peshawar 25120, Khyber Pakhtunkhwa, Pakistan


³ Center for Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas (UFPEL), Pelotas, Brazil

⁴ Department of Biochemistry, University of Malakand, Chakdara Dir Lower, Totakan 18800, Khyber Pakhtunkhwa, Pakistan

⁵ Department of Chemistry, University of Liverpool, Liverpool, UK

⁶ Department of Pharmacognosy, College of Pharmacy, King Saud University, 11451 Riyadh, Saudi Arabia

⁷ Department of Surgery, College of Medicine, King Saud University, P.O.Box 7805, 11472 Riyadh, Kingdom of Saudi Arabia



Advances in liposomes loaded with photoresponse materials for cancer therapy

Xianwei Zhang^a, Youfu Ma^a, Yenong Shi^a, Lihe Jiang^b, Lisheng Wang^c, Haroon ur Rashid^c, Mingqing Yuan^{a,*}, Xu Liu^{a,*}

^a Guangxi Key Laboratory of Special Biomedicine, School of Medicine, Guangxi University, Nanning 530004, China

^b School of Basic Medical Sciences, Youjiang Medical University for Nationalities, Baise, Guangxi 533000, China

^c Center for Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas (UFPEL), Pelotas, RS 96010-900, Brazil

ARTICLE INFO

Keywords:

Liposomes
Phototherapy
Photoresponsive nanomaterials
Delivery system
Synergistic therapy

ABSTRACT

Cancer treatment is presently a significant challenge in the medical domain, wherein the primary modalities of intervention include chemotherapy, radiation therapy and surgery. However, these therapeutic modalities carry side effects. Photothermal therapy (PTT) and photodynamic therapy (PDT) have emerged as promising modalities for the treatment of tumors in recent years. Phototherapy is a therapeutic approach that involves the exposure of materials to specific wavelengths of light, which can subsequently be converted into either heat or Reactive Oxygen Species (ROS) to effectively eradicate cancer cells. Due to the hydrophobicity and lack of targeting of many photoresponsive materials, the use of nano-carriers for their transportation has been extensively explored. Among these nanocarriers, liposomes have been identified as an effective drug delivery system due to their controllability and availability in the biomedical field. By binding photoresponsive materials to liposomes, it is possible to reduce the cytotoxicity of the material and regulate drug release and accumulation at the tumor site. This article provides a comprehensive review of the progress made in cancer therapy using photoresponsive materials loaded onto liposomes. Additionally, the article discusses the potential synergistic treatment through the combination of phototherapy with chemo/immuno/gene therapy using liposomes.

1. Introduction

Cancer remains a major health crisis affecting mankind. Over the next 20 years, the number of cancer cases worldwide is projected to increase by approximately 50% [1]. While chemotherapy (CT), radiation therapy (RT) and surgical treatment are commonly used to treat cancer, these methods often come with side effects. Long-term monotherapy with chemotherapy drugs is often limited by therapeutic efficacy, drug resistance, metastasis, and adverse side effects [2]. Radiation therapy can damage normal cells, leading to side effects such as

secondary cancers, blisters, and other skin problems [3–5]. Surgical treatment is both risky and traumatic for the patient, and it is also prone to complications [6]. However, due to the unclear etiology of certain tumors, it is difficult to cure completely. Moreover, the proliferation of malignant tumors can lead to disturbances in normal physiological functions of the body, such as the inability of the immune system to initiate normal immune processes to suppress tumor growth. Tumor drug resistance is perhaps the greatest challenge in the treatment of late-stage cancer. The reasons leading to tumor drug resistance can be divided into the following aspect: tumor heterogeneity, undruggable

Abbreviations: CT, chemotherapy; RT, radiation therapy; PTT, photothermal therapy; PDT, photodynamic therapy; EPR, enhanced permeability and retention; PTCAs, photothermal conversion agents; PEG, polyethylene glycol; UV, unilamellar vesicles; LUV, large unilamellar vesicles; GUV, giant unilamellar vesicles; MLV, multilamellar vesicles; ROS, reactive oxygen species; LDDS, Liposomal drug delivery systems; VEGF, vascular endothelial growth factor; NIR, Near Infrared; PTAs, photothermal agents; ICG, Indocyanine green; PA, photoacoustic; LSPR, local surface plasmon resonance; NPs, nanoparticles; GA, gambogic acid; HSP90, heat shock protein 90; CNTs, carbon nanotubes; GO, graphene oxide; CDs, carbon dots; RGO, graphene oxide.; DOX, Doxorubicin; PL, peptide lipid; SL, sucrose laurate; siRNAs, small interfering RNAs; PSs, photosensitizers; ACQ, aggregation-induced quenching; BP, bis-pyrene; TME, tumor microenvironment; CAT, catalase; GSH, glutathione; TAMs, Tumor-associated macrophages; TAAs, tumor-associated antigens; IERL, immune-enhancing liposome; UCNPs, upconversion nanoparticles; ITH, intratumor heterogeneity; ILPCR, immunoliposome polymerase chain reaction; ILPCR, immunoliposome polymerase chain reaction.; CSCs, cancer stem cells.

* Corresponding authors.

E-mail addresses: yuanmingqing1985@163.com (M. Yuan), wendaoliuxu@163.com (X. Liu).


<https://doi.org/10.1016/j.bioph.2024.116586>

Received 22 February 2024; Received in revised form 7 April 2024; Accepted 10 April 2024

0753-3322/© 2024 The Authors. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

^{31}P Nuclear Magnetic Resonance Spectroscopy for Monitoring Organic Reactions and Organic Compounds

João Marcos Anghinoni, Irum, **Haroon Ur Rashid**, Eder João Lenardão,* and Márcio Santos Silva*[a]



Research Article


Muhammad Kashif, Sumaira Naz, Muhammad Zahoor*, Syed Wadood Ali Shah, Jalal Uddin, Muhammad Esa, Haroon ur Rashid, Riaz Ullah, Amal Alotaibi

Antioxidant and antidiabetic potentials of methoxy-substituted Schiff bases using *in vitro*, *in vivo*, and molecular simulation approaches

<https://doi.org/10.1515/chem-2024-0083>

received June 1, 2024; accepted August 27, 2024

Abstract: The current study attempted to synthesize methoxy-substituted Schiff's bases, namely MK1 and MK2, and evaluate their antidiabetic effects using *in vitro*, *in vivo*, and molecular docking studies. Experimental animals (rat model) received the synthetic compounds, MK1 and MK2, orally in doses of 25 and 50 mg/kg body weight, respectively. When comparing compound MK2 at the tested doses to glibenclamide on day 28, the diabetic rats' blood glucose levels were nearly normal (139.02 and 121.23 mg/dL at 25 and 50 mg/kg body weight doses). The IC_{50} for MK1 against α -glucosidase inhibitory potential was found to be 281.29 μ g/mL, while for MK2, it is reported to be 204.69 μ g/mL. Furthermore, the acute toxicity, lipid profile, and its effect on blood biochemical parameters were also examined. In addition, through *in silico* analysis, the binding of MK1



Graphical abstract

and MK2 was elucidated with α -glucosidase enzyme, showcasing its antidiabetic mechanism at molecular levels. The *in silico* studies also predicted the two compounds to be inactive toward the human hERGs cardiac potassium channel, which indicates no potential risk of cardiac toxicity. Overall, the toxicity predictions suggest that compounds MK1 and MK2 are non-toxic and non-carcinogenic.

Keywords: Schiff's bases, antidiabetic potential, diabetes, glibenclamide

1 Introduction

Diabetes mellitus (DM), one of the metabolic illnesses, is characterized by a persistent increase in the level of blood glucose [1]. This is associated with the interruption of the metabolism of carbohydrates, proteins, and lipids. This variation in metabolism arises due to the low amount of insulin production from the pancreas or because the body does not use the insulin produced [2]. It ranks among the top ten causes of adult mortality, accounting for an estimated four million deaths globally in 2023 [3]. The biomarkers of diabetes include fasting blood sugar, hemoglobin A1c (HbA1c), insulin, β -cell function markers (e.g., HOMA-B), triglycerides (TG), lipid profile, and inflammatory

* **Corresponding author: Muhammad Zahoor**, Department of Biochemistry, University of Malakand, Chakdara, 18800, Pakistan, e-mail: mohammadzahoorus@yahoo.com

Muhammad Kashif: Department of Biochemistry, University of Malakand, Chakdara, 18800, Pakistan, e-mail: mkashif00447@gmail.com

Sumaira Naz: Department of Biochemistry, University of Malakand, Chakdara, 18800, Pakistan, e-mail: sumaira@uom.edu.pk

Syed Wadood Ali Shah: Department of Pharmacy, University of Malakand, Chakdara, 18800, Khyber Pakhtunkhwa, Pakistan, e-mail: pharmacistsyed@gmail.com

Jalal Uddin: Department of Biochemistry, University of Malakand, Chakdara, 18800, Pakistan, e-mail: jalalchm@gmail.com

Muhammad Esa: Department of Pharmacy, Shaheed Benazir Bhutto University, Sheringal, Dir Upper, Pakistan, e-mail: esakhan5595@gmail.com

Haroon ur Rashid: Center of Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, RS, Brazil, e-mail: haroongold@gmail.com

Riaz Ullah: Medicinal Aromatic and Poisonous Plants Research Center, Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, 11451, Saudi Arabia, e-mail: rullah@ksa.edu.sa

Amal Alotaibi: Department of Basic Science, College of Medicine, Princess Nourah bint Abdulrahman University, Riyadh, 11671, Saudi Arabia, e-mail: amaalotaibi@pnu.edu.sa

Research Article

Gulab Said, Waqas Ahmad, Muhammad Tariq Jan, Muhammad Zahoor*, Azmat Ullah, Riaz Ullah, Muhammad Naveed Umar, Zafar Iqbal and Haroon ur Rashid

Removal of metaldehyde pesticide from aquatic media using modified cellulose obtained from *Populus nigra* plant, as potential adsorbent

<https://doi.org/10.1515/pac-2024-0234>

Abstract: In this study modified cellulose based adsorbent was prepared from *Populus nigra* plant, and used for elimination of metaldehyde (herbicide) from model waste water. The adsorbent was characterized through analytical technique such as FTIR, SEM, EDX and XRD for structural adsorption related parameters. The results of SEM showed the suitability of the material to be used as adsorbent and FTIR showed successful crosslinking of polyvinyl alcohol into cellulose structure. In order to get maximum reclamation benefits from adsorbent it was subjected to a number of tests evaluating the effect of metaldehyde concentration, sorbent dose, contact time, initial pH of solution and temperature. The maximum removal of 70 % was achieved under conditions of 80 mg/L metaldehyde concentration, 60 min contact time, pH of 8, 0.08 g sorbent dosage, and room temperature (25 °C). The Langmuir isotherm model with correlation coefficients of 0.9855 and maximum adsorption capacity recorded was 8.32 mg/g, while excellent agreement was shown by kinetic data with pseudo second order kinetic model with $R^2 = 0.9876$. Thermodynamic study indicated enthalpy change ($\Delta H^\circ = -129$ kJ/mol) to be negative, entropy change ($\Delta S^\circ = 161.7$ J/mol) positive, and the Gibbs free energy (ΔG) as negative showing that the process to be exothermic and feasible/spontaneous with an increase of randomness at solid liquid interface. The finding indicated that modified cellulose could be used as an efficient adsorbent for removal of metaldehyde from model waste water. However, further validation with other pollutants will be helpful in checking reproducibility of the present findings.

Keywords: Acetylated cellulose; isothermal study; kinetic study; metaldehyde; polyvinyl alcohol; *Populus nigra*.

Introduction

The increase in demand of food due to increasing world population have led to increased agriculture activities which in turn have increased the use of herbicides globally. There are different types of herbicides used to increase crops production. However, many of these herbicides are persistent and remains intact in different

***Corresponding author: Muhammad Zahoor**, Department of Biochemistry, University of Malakand, Chakdara, Dir Lower, Pakistan, e-mail: mohammadzahoorus@yahoo.com. <https://orcid.org/0000-0002-4528-8517>

Gulab Said and Muhammad Tariq Jan, Department of Chemistry, Islamia College Peshawar, Peshawar 25120, Pakistan, e-mail: gulab_said@yahoo.com (G. Said), tariq@icp.edu.pk (M.T. Jan)

Waqas Ahmad, Department of Chemistry, ICS, University of Peshawar, Peshawar, 25121, Pakistan, e-mail: waqasahmad@uop.edu.pk

Azmat Ullah, Department of Chemistry Government Degree College, Mingora Swat, 19120, Pakistan, e-mail: azmatullah9499@gmail.com

Riaz Ullah, Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia, e-mail: rullah@ksu.edu.sa

Muhammad Naveed Umar, Department of Chemistry, University of Liverpool, Liverpool, UK, e-mail: m.naveed-umar@liverpool.ac.uk

Zafar Iqbal, Department of Surgery, College of Medicine, King Saud University, P.O. Box 7805, Riyadh, 11472, Kingdom of Saudi Arabia, e-mail: zsyed@ksu.edu.sa

Haroon ur Rashid, Center of Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, RS, Brazil, e-mail: haroongold@gmail.com. <https://orcid.org/0000-0003-3335-7957>

Research Article


Nasir Ahmad, Khalid Khan, Sher Wali Khan, **Haroon Ur Rashid***, Irum, Muhammad Zahoor*, Muhammad Naveed Umar, Riaz Ullah, Essam A. Ali

Homology modeling and molecular docking study of corticotrophin-releasing hormone: An approach to treat stress-related diseases

<https://doi.org/10.1515/chem-2024-0069>

received February 22, 2024; accepted June 30, 2024

Abstract: Corticotropin-releasing hormone receptors (CRHRs), also termed corticotropin-releasing factor receptors, are linked to G-protein-coupled receptor class. Corticotropin-releasing hormone (CRH) is medically significant in stress, immune response, gastrointestinal motility, and eating patterns. It serves as a releasing hormone and is encoded by the CRH gene. It has been established that there are two subtypes of CRHRs: CRH1-R and CRH2-R. These receptors, representing types 1 and 2, respectively, play a crucial role in regulating biological functions triggered by CRH. To treat stress-related gut abnormalities and stress-related disorders, regulation and optimization of CRH1-R and CRH2-R have turned into a novel idea. The three-dimensional (3D) structure of CRH is not completely recognized, and it is believed that the peptide key unit is helical and both the ultimate edges are relatively unsaturated. We can envisage its 3D structure from the amino acid order of a model protein by homology modeling procedures using Molecular Operating Environment and the Iterative Threading Assembly Refinement program. The assessment



Graphical abstract

and authentication of the 3D structure were performed with RAMPAGE and ERRATE online servers. Utilizing the 3D structure of the target protein and predictions of its active site assists us in the development of new drug candidates aimed at treating disorders associated with stress. CRHR was docked with 19 CP376395 analogs acting as antagonists.

Keywords: homology modeling, molecular docking, corticotrophin, adenylate cyclase, nucleotides

1 Introduction

G-protein-coupled receptors (GPCRs) facilitate most of our biological feedback to neurotransmitters, hormones, and environmental stimulants, and so they possess a significant potential to act as therapeutic targets for a variety of diseases. [1]. The structure of GPCRs is usually comprised of seven α -helical transmembrane (TM) domains, an intracellular C-terminus, and an extracellular N-terminus [2,3]. Hydrophobic seven TM helical central core domain is linked by three intracellular and three extracellular loops. In extracellular loops 1 and 2, two cysteine residues are conserved to form a disulfide bridge, which is important for the stabilization and packing of a restricted number of conformations of seven TMs [4,5]. They are included in the cell membranes of a wide range of organisms including invertebrates, microorganisms, plants, and mammals [6,7]. All types of GPCRs share a common membrane topology, the

* **Corresponding author: Haroon Ur Rashid**, Center of Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, 96010-900, RS, Brazil, e-mail: haroongold@gmail.com

* **Corresponding author: Muhammad Zahoor**, Department of Biochemistry, University of Malakand, Chakdara Dir Lower, 18800, Khyber Pakhtunkhwa, Pakistan, e-mail: mohammadzahoorus@yahoo.com

Nasir Ahmad, Khalid Khan: Department of Chemistry, Islamia College University, Peshawar, 25120, Khyber Pakhtunkhwa, Pakistan

Sher Wali Khan: Department of Chemistry, Rawalpindi Women University, 6th Road Satellite Town, Rawalpindi, 46300, Punjab, Pakistan

Irum: Center of Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, 96010-900, RS, Brazil

Muhammad Naveed Umar: Department of Chemistry, University of Liverpool, Liverpool, United Kingdom

Riaz Ullah: Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, 11451, Saudi Arabia

Essam A. Ali: Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia

Nasir Ahmad, Khalid Khan*, Haroon ur Rashid*, Sher Wali Khan, Muhammad Naveed Umar, Naila Gulfam, Muhammad Zahoor*, Riaz Ullah and Essam A. Ali

Homology modeling and molecular docking study of metabotropic glutamate receptor 5 variant F: an attempt to develop drugs for treating CNS diseases

<https://doi.org/10.1515/zpch-2023-0449>

Received November 7, 2023; accepted January 9, 2024; published online January 23, 2024

Abstract: G protein-coupled receptors (GPCRs) constitute the largest protein family involved in signal transduction and are the main targets for drug delivery. The metabotropic glutamate receptors (mGluRs) are class C of GPCRs that contribute to the modulation of glutamatergic neurotransmission which regulates vital physiological processes and is recognized as a potential therapeutic target throughout the central nervous system. Intracellular C-terminal domains of mGluR are the main targets for proteins regulating these receptors and are also involved in alternative splicing, regulation by phosphorylation, and modulatory protein-protein interactions. Class C of GPCRs exist as dimers and is mediated by interactions

***Corresponding authors:** Khalid Khan, Department of Chemistry, Islamia College University Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan, E-mail: drkhalidchem@yahoo.com; Haroon ur Rashid, Center of Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, RS, Brazil, E-mail: haroongold@gmail.com; and Muhammad Zahoor, Department of Biochemistry, University of Malakand, Chakdara Dir Lower, Khyber Pakhtunkhwa, 18800, Pakistan, E-mail: mohammadzahoorus@yahoo.com

Nasir Ahmad, Department of Chemistry, Islamia College University Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan, E-mail: nasirdft@gmail.com

Sher Wali Khan, Department of Chemistry, Rawalpindi Women University, 6th Road Satellite Town, Rawalpindi, Pakistan, E-mail: sherwalikhn@yahoo.co.uk

Muhammad Naveed Umar, Department of Chemistry, University of Liverpool, Liverpool, UK; and Department of Chemistry, University of Malakand, Chakdara Dir Lower, Khyber Pakhtunkhwa, 18800, Pakistan, E-mail: m.naveed-umar@liverpool.ac.uk

Naila Gulfam, Department of Zoology, Jinnah College for Women, University of Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan, E-mail: nailazoo@uop.edu.pk

Riaz Ullah, Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, E-mail: rullah@ksu.edu.sa

Essam A. Ali, Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia, E-mail: esali@ksu.edu.sa

ARTICLE

Molecular Mechanisms of Buzhong Yiqi Decoction in the Treatment of Gastric Cancer: A Network Pharmacology, Molecular Docking, and In Vitro Experimental Analysis

Panke Zeng¹  | Xinyu Wu² | Chen Chen³ | Jianing Zhang¹ | Haroon ur Rashid⁴ | Pengfei Zhang²

¹Department of Pharmacy, The First Affiliated Hospital, Xi'an Jiaotong University, Xi'an, China | ²The First Clinical Medical College, Shandong University of Traditional Chinese Medicine, Jinan, China | ³Department of Pharmacy, Taihe Hospital, Shiyao, China | ⁴Center for Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, RS, Brazil

Correspondence: Panke Zeng (j_hope_medicine@xjtu.edu.cn) | Pengfei Zhang (65090753@sdu.edu.cn)

Received: 13 April 2024 | **Revised:** 12 June 2024 | **Accepted:** 14 June 2024

Funding: This work was supported by the Key Research and Development Program of Shaanxi (2024SF-YBXM-111).

Keywords: Buzhong Yiqi decoction | experimental verification | gastric cancer | molecular docking | molecular mechanisms | network pharmacology

ABSTRACT

Gastric cancer (GC) is the most common type of cancer of the digestive system with high morbidity and mortality. Chemotherapy and targeted therapy are used to treat patients with advanced GC. However, side effects and drug resistance to the two modalities remain the main challenges. The Buzhong Yiqi decoction (BZYQD), a classical traditional Chinese medicine formula, has been reported for the treatment of various types of cancers. However, the underlying pharmacological mechanism has not been fully elucidated. Therefore, this study integrated network pharmacology, molecular docking, cancer public databases, and cell experiments to explore the potential bioactive compounds and BZYQD's mechanism of action against GC. A total of 245 targets of BZYQD, 5291 GC-related targets, and 186 were identified as their common targets through the database. Network analysis confirmed AKT1, TP53, TNF, and EGFR to be the core targets, while the main compounds observed were quercetin, kaempferol, and β -Sitosterol. The core signaling pathways included PI3K-AKT, MAPK, TNF, and IL-17. Molecular docking revealed good binding activity for the main compounds and core targets. Based on the database's validation of core targets, a large number of core genes were verified to be consistent with this study. Quercetin, kaempferol, and β -Sitosterol were found to significantly reduce the growth of GC cells in the MTT experiment. The current study revealed that BZYQD may inhibit GC progression by interfering with core targets such as AKT1, TP53, TNF, EGFR, and MAPK3, and by regulating the activity of PI3K-AKT, MAPK, TNF, and IL-17 signaling pathways.

1 | Introduction

Gastric cancer (GC) is one of the most prevalent malignancies and the world's fourth-leading cause of cancer-related deaths. The occurrence and progression of GC are influenced by several factors, including family genetic history, irregular diet, excessive alcohol consumption, and *Helicobacter pylori*

infection. Current statistics indicate that morbidity and mortality rates are elevated in many countries [1]. Additionally, GC stands as the second-leading cause of cancer morbidity and mortality in China. In comparison to the developed countries, China exhibits higher 5-year incidence (27.6/100,000) and mortality/incidence ratio (0.845) rates. Furthermore, the incidence of early-onset gastric cancer (EOGC) in the young

Panke Zeng and Xinyu Wu shared the first authorship.

© 2024 Wiley Periodicals LLC.

REVIEW ARTICLE

Recent Advances on the Antimicrobial Activities of Schiff Bases and their Metal Complexes: An Updated Overview

Juliana Jorge^{1,#}, Kristiane Fanti Del Pino Santos^{1,#}, Fernanda Timóteo¹, Rafael Rodrigo Piva Vasconcelos¹, Osmar Ignacio Ayala Cáceres¹, Isis Juliane Arantes Granja², David Monteiro de Souza Jr.¹, Tiago Elias Allievi Frizon³, Giancarlo Di Vaccari Botteselle⁴, Antonio Luiz Braga⁵, Sumbal Saba^{2,*}, Haroon ur Rashid^{1,5,*} and Jamal Rafique^{1,2,*}

¹Instituto de Química, Universidade Federal do Mato Grosso do Sul, Campo Grande, 79074-460, MS, Brazil;

²Instituto de Química, Universidade Federal de Goiás - UFG, Goiânia, 74690-900, GO, Brazil; ³Department of Energy and Sustainability, Universidade Federal de Santa Catarina - UFSC, Campus Araranguá, Araranguá, 88905-120, SC, Brazil; ⁴Departamento de Química, Universidade Estadual do Centro-Oeste - UNICENTRO, Guarapuava, 85819110, PR, Brazil; ⁵Departamento de Química, Universidade Federal de Santa Catarina, 88040-970, Florianópolis, SC, Brazil

ARTICLE HISTORY

Received: August 03, 2022
Revised: December 12, 2022
Accepted: December 29, 2022

DOI:
10.2174/0929867330666230224092830



Abstract: Schiff bases represent a valuable class of organic compounds, synthesized via condensation of primary amines with ketones or aldehydes. They are renowned for possessing innumerable applications in agricultural chemistry, organic synthesis, chemical and biological sensing, coating, polymer and resin industries, catalysis, coordination chemistry, and drug designing. Schiff bases contain imine or azomethine ($-C=N-$) functional groups which are important pharmacophores for the design and synthesis of lead bioactive compounds. In medicinal chemistry, Schiff bases have attracted immense attention due to their diverse biological activities. This review aims to encompass the recent developments on the antimicrobial activities of Schiff bases. The article summarizes the antibacterial, antifungal, antiviral, antimalarial, and antileishmanial activities of Schiff bases reported since 2011.

Keywords: Schiff base, antimicrobial activity, imine, azomethine, antibacterial, antifungal, antiviral, antimalarial.

1. INTRODUCTION

Schiff bases are a class of nitrogenous organic compounds that have a general structure $R-CH=N-R'$, where R and R' can be an alkyl or aryl group, linear or cyclic, which can be substituted in many ways [1-3]. They can be considered a subclass of imines, being secondary ketamines (when both R and R' are different from H) and secondary aldimines (when R or R' is an H atom), depending on the presented structure [1, 2, 4, 5].

In the literature, they are considered synonyms for imines ($>C=N-$) and azomethines (or secondary aldimines, $-HC=N-$), as they are also known [1, 4-8]. They received this name because of Hugo Schiff (1834-1915), an Italian-German chemist laureated with the Nobel Prize and brother of the physiologist Moritz Schiff [1, 2, 4]. Hugo Schiff also discovered other imines, in addition to studying aldehydes, which led to the development of the Schiff reagent, for the identification of these compounds, which consists of a colorless solution containing an indicator produced by the reaction between pararosaniline with sulfite in an acid medium [5, 6].

The formation reaction of a Schiff base is presented according to the Schiff methodology (Fig. 1). They can be synthesized from an aliphatic or aromatic primary amine (of the type $R-NH_2$ or $Ar-NH_2$), with an active

*Address correspondence to these authors at the Instituto de Química, Universidade Federal de Goiás - UFG, Goiânia, 74690-900, GO, Brazil; E-mail: sumbalsaba@ufg.br (S.S.); Departamento de Química, Universidade Federal de Santa Catarina, 88040-970, Florianópolis, SC, Brazil; E-mail: haroongold@gmail.com (H.R.); Instituto de Química, Universidade Federal do Mato Grosso do Sul, Campo Grande, 79074-460, MS, Brazil; E-mails: jamal.rafique@ufms.br, jamal.chm@gmail.com (J.R.)

[#]Both authors contributed equally to this work.

RESEARCH ARTICLE

Integrating network pharmacology and experimental verification to investigate the pharmacological mechanisms of Buzhong Yiqi decoction in the treatment of non-small cell lung cancer

Panke Zeng¹  | Feng Wang¹ | Jianing Zhang¹ | Haroon ur Rashid² | Xin Li¹ | Pengfei Zhang³ | Yunru Luo³ | Xinyu Wu³

¹Department of Pharmacy, The First Affiliated Hospital, Xi'an Jiaotong University, Xi'an, China

²Center for Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas (UFPEL), Pelotas, Brazil

³The First Clinical Medical College, Shandong University of Traditional Chinese Medicine, Jinan, China

Correspondence

Panke Zeng, Department of Pharmacy, The First Affiliated Hospital, Xi'an Jiaotong University, Xi'an 710061, China.

Email: j_hope_medicine@xjtuqh.edu.cn

Xinyu Wu, The First Clinical Medical College, Shandong University of Traditional Chinese Medicine, Jinan 250355, China.

Email: wxysgxa666@163.com

Abstract

Among all types of cancers, non-small cell lung cancer (NSCLC) exhibits the highest mortality rate with a five-year survival rate below 17% for patients. The Buzhong Yiqi decoction (BZYQD), traditional Chinese medicine (TCM) formula, has been reported to exhibit clinical efficacy in the treatment of NSCLC. Nevertheless, the underlying molecular mechanism remains elusive. This study aimed to assess the mechanistic actions exerted by BZYQD against NSCLC using network pharmacological analysis and experimental validation. The public databases were searched for active compounds in BZYQD, their potential targets, and NSCLC-related targets. The protein–protein interaction (PPI) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses were performed to predict the core targets and signaling pathways of BZYQD against NSCLC. After screening, this study validated the results of predictions through in vitro experiments and public databases. We found 192 common targets between BZYQD and NSCLC. KEGG analysis showed that the anti-NSCLC effects of BZYQD were mediated through the PI3K-AKT signaling pathway. The results of in vitro experiment indicated that BZYQD could inhibit cell viability and proliferation of A549 and H1299 cells apart from inducing cell apoptosis. In addition, western blot results substantiated that BZYQD could treat NSCLC by inhibiting the activation of the PI3K-AKT signaling pathway. The current study investigated the pharmacological mechanism of BZYQD against NSCLC via network pharmacology and in vitro analyses. Overall, the results revealed that BZYQD could be a promising therapeutic agent for the treatment of NSCLC in the future. Still, more experimental investigations are needed to confirm the applicability of BZYQD for clinical trials.

KEYWORDS

Buzhong Yiqi decoction, experimental verification, network pharmacology, non-small cell lung cancer, PI3K-AKT signaling pathway

Homology Modeling of Alpha-Glucosidase from *Candida albicans*: Sequence Analysis and Structural Validation Studies *in silico*

Haroon Ur Rashid,^{a,b} Vanderlan S. Bolzani,^a Khalid Khan,^c Luiz Antonio Dutra,^a
Nasir Ahmad^{*,c} and Abdul Wadood^{*,d}

^aInstituto de Química, Universidade Estadual Paulista Julio de Mesquita Filho,
14800-060 Araraquara-SP, Brazil

^bCentro de Ciências Químicas, Farmacêuticas e de Alimentos, Universidade Federal de Pelotas,
96010-900 Pelotas-RS, Brazil

^cDepartment of Chemistry, Islamia College University, Peshawar, 25120, Khyber Pakhtunkhwa,
Pakistan

^dDepartment of Biochemistry, Shankar Campus, Abdul Wali Khan University, Mardan, 23200,
Khyber Pakhtunkhwa, Pakistan

The alpha-glucosidase enzyme of *Candida albicans* plays a vital role in the pathogenesis of candidiasis, a serious fatal disease in immune-compromised patients. The unavailability of the three-dimensional crystallographic structure of this enzyme creates a hindrance in developing novel and potent inhibitors. Here, an attempt has been made to design a stable three-dimensional conformer of alpha-glucosidase through *in silico* analysis which may be helpful for the designing of effective drugs. For this purpose, the oligo-1-6-glucosidase enzyme is used as a template for homology modeling of the alpha-glucosidase structure by Molecular Operating Environment 2011-12 software. The generated model was validated through ERRAT and Ramachandran tools, whereas its stability was studied through the molecular dynamics simulation technique. The obtained results indicate that model of alpha-glucosidase has stable secondary and tertiary arrangements. This finding may spur new directions for the rational designing and development of new antifungal inhibitors. Nevertheless, additional experimental investigations and validation are needed to confirm the *in silico* results of this study.

Keywords: alpha-glucosidase, candidiasis, homology modeling, simulations

Introduction

The infections associated with the fungus have been enhanced at a large scale which leads to the enhancement of the mortality rate in humans. The species of *Candida* are mostly human pathogens responsible for infections of deep tissues and mucosa. The species of candida are related to the microbiota of a patient mucosal oral cavity vagina and gastrointestinal tract;¹ and cause different diseases ranging from an overgrowth of mucocutaneous to the infections of blood² which include candidemia a major disease that causes death. Candidiasis is included among the infections of *Candida* species.³ *Candida* infection may present as

angular cheilitis, acute pseudomembranous, acute atrophic, chronic atrophic, or chronic hyperplastic candidiasis. There are many reasons like diabetes, malignancies, and weak immunity which strengthen oropharyngeal candidiasis. Denture stomatitis and oropharyngeal candidiasis are caused by *Candida albicans*. The occult esophageal infection and oral fungal disease are linked with each other while the past study of odynophagia warned medical scientists about the development of systemic candidiasis.⁴ Normally, healthy humans also carry the above-mentioned fungus but they caused disease in human immunodeficiency virus (HIV) patients because they have weak immunity. The composition of the genus involved a diverse group that included 17 different *Candida* species which cause infection in humans about 90% of infections are caused by *Candida albicans*, *Candida glabrata*, *Candida parapsilosis*,

*e-mail: nasirdft@gmail.com, awadood@awkum.edu.pk
Editor handled this article: Paula Homem-de-Mello (Associate)



Research Article

Sher Wali Khan*, Saira Nayab, Muhammad Naveed Umar, Momin Khan, Anila Iqbal, Nasir Ahmad, Haroon ur Rashid, Muhammad Ishaq Ali Shah, Naila Gulfam, Muhammad Zahoor*, Riaz Ullah, Essam A. Ali

Synthesis, characterization, *in silico* and *in vitro* studies of novel glycoconjugates as potential antibacterial, antifungal, and antileishmanial agents

<https://doi.org/10.1515/chem-2023-0195>

received November 11, 2023; accepted January 12, 2024

Abstract: In the present work, several new glycoconjugates (**8a–e**) were generated from glycopyranosyl- α -trichloroacetimidates (sugar-OTCA) as glycosyl donors and dimethyl-L-

tartrate as an aglycone acceptor in good to excellent yields. In the synthetic protocol, various monosaccharides were transformed into pentaacetylated derivatives and then into glycopyranosyl- α -trichloroacetimidates. Afterward, the sugar-OTCA was reacted with dimethyl-L-tartrate using Schmidt's trichloroacetimidate protocol to give the desired products. The newly synthesized glycoconjugates were characterized by FT-IR, ^1H , and ^{13}C -NMR spectroscopic analytical methods. All the target compounds (**8a–e**) were tested *in vitro* against various strains of bacteria and fungi at different concentrations. The results revealed that the target compounds had encouraging antibacterial and antifungal potential. The antileishmanial activity of the target compounds against *Leishmania tropica* promastigotes was also investigated. The *in vitro* results were further supported by the *in silico* docking study that indicated minimum values of the docking scores and binding energies for the resulting complexes obtained by the favorable interactions between the target compounds (**8a–e**) and the selected strains of bacteria and fungi. The docking results proposed promising antibacterial and antifungal activities of the target compounds (**8a–e**) against the selected bacterial and fungal species.

Keywords: glycoconjugates, glycopyranosyl- α -trichloroacetimidates, glycosylation, antimicrobial, antileishmanial, molecular docking

* **Corresponding author: Sher Wali Khan**, Department of Chemistry, The Rawalpindi Women University, 6th Road Satellite Town, Rawalpindi, Pakistan, e-mail: sherwalikhan@f.rwu.edu.pk

* **Corresponding author: Muhammad Zahoor**, Department of Biochemistry, University of Malakand, Chakdara Dir Lower, KPK, 18800, Pakistan, e-mail: mohammadzahoorus@yahoo.com

Saira Nayab: Department of Chemistry, Shaheed Benazir Bhutto University Sheringal, Dir (Upper), Khyber Pakhtunkhwa, Pakistan; Department of Chemistry and Green-Nano Materials Research Center, Kyungpook National University, Daegu, 41566, Republic of Korea

Muhammad Naveed Umar: Department of Chemistry, University of Liverpool, Liverpool, United Kingdom; Department of Chemistry, University of Malakand, Chakdara Dir Lower, KPK, 181800, Pakistan, e-mail: m.naveed-umar@liverpool.ac.uk

Momin Khan: Institute of Pathology and Diagnostic Medicines (IPDM), Khyber Medical University, Peshawar, KP, Pakistan, e-mail: mominkhan.ibms@kmu.edu.pk

Anila Iqbal: National Center for Physics (NCP), Quaid-i-Azam University, Islamabad, 45320, Pakistan, e-mail: a.anilaiqbal@gmail.com

Nasir Ahmad: Department of Chemistry, Islamia College University Peshawar, Khyber Pakhtunkhwa, Pakistan, e-mail: nasirdft@gmail.com

Haroon ur Rashid: Center for Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas, RS, Brazil, e-mail: haroongold@gmail.com

Muhammad Ishaq Ali Shah: Department of Chemistry, Abdul Wali Khan University, Mardan, KP, Pakistan, e-mail: ishaqalishah@awku.edu.pk

Naila Gulfam: Department of Zoology, Jinnah College for Women, University of Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan, e-mail: nailazoo@uop.edu.pk

Riaz Ullah: Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, e-mail: rullah@ksu.edu.sa

Essam A. Ali: Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia, e-mail: esali@ksu.edu.sa

1 Introduction

Carbohydrates (biomolecules) are often found attached to other biomolecules that constitute glycoconjugates. Glycolipids and glycoproteins are normally found on cell surfaces that act as receptors for cell–cell determination and are involved in cell growth, repair adhesion, migration, and tumor metastasis in cancer [1]. Vaccination is one of the most cost-effective ways and a key strategy for controlling various infectious diseases



OPEN

Biological investigations of *Aspergillus ficuum* via in vivo, in vitro and in silico analyses

Zafar Ali Shah¹, Khalid Khan^{1✉}, Tanzeel Shah², Nasir Ahmad¹, Akhtar Muhammad¹ & Haroon ur Rashid^{3,4✉}

Serious human health impacts have been observed worldwide due to several life-threatening diseases such as cancer, candidiasis, hepatic coma, and gastritis etc. Exploration of nature for the treatment of such fatal diseases is an area of immense interest for the scientific community. Based on this idea, the genus *Aspergillus* was selected to discover its hidden therapeutic potential. The genus *Aspergillus* is known to possess several biologically active compounds. The current research aimed to assess the biological and pharmacological potency of the extracts of less-studied *Aspergillus ficuum* (FCBP-DNA-1266) (*A. ficuum*) employing experimental and bioinformatics approaches. The disc diffusion method was used for the antifungal investigation, and the MTT assay was performed to assess the anticancer effects. Mice were employed as an in vivo model to evaluate the antispasmodic effects. A standard spectrophotometric technique was applied to gauge the urease inhibitory activity. The antifungal studies indicate that both n-hexane and ethyl acetate extracts were significantly active against *Candida albicans* (*C. albicans*) with their zone of inhibitions (ZOI) values reported as 19 ± 1.06 mm and 25 ± 0.55 mm, respectively at a dose of $30 \mu\text{g.mL}^{-1}$. In vitro cytotoxicity assay against HeLa, fibroblast 3T3, prostate PC3, and breast MCF-7 cancer cell lines was performed. The ethyl acetate extract of *A. ficuum* was found to be significantly active against MCF-7 with its IC_{50} value of $43.88 \mu\text{g.mL}^{-1}$. However, no substantial effects on the percent cell death of HeLa cancer cell lines were observed. In addition, the *A. ficuum* extracts also inhibited the urease enzyme compared to standard thiourea. The antispasmodic activity of *A. ficuum* extract was assessed by an in vivo model and the results demonstrated promising activity at 150 mg.kg^{-1} . Molecular docking results also supported the antifungal, anticancer, and antiurease potency of *A. ficuum* extract. Overall, the results display promising aspects of *A. ficuum* extract as a future pharmacological source.

New drugs are needed due to certain factors such as globalization, the aging population, and microbial resistance to existing medicines. Only a small number of biologically active compounds are currently used in prescription drugs¹. Scientists are currently investigating emerging diseases to gain a better understanding of them and find their potential cures using various natural and chemical formulations. However, there are still many areas that remain unexplored due to insufficient knowledge and techniques. To find new bioactive compounds that can be used for medicinal, agricultural, and industrial purposes, researchers are focusing on isolating such compounds from new species of fungi².

In recent times, researchers have isolated several metabolites from fungi, including anthraquinones, alkaloids, terpenes, steroids, flavonoids, and cyclic peptides. Some secondary metabolites produced by various fungi have been found to exhibit potent biological activities such as antifungal, antitumor, anti-inflammatory antibacterial, antiparasitic, and antiviral properties. The chemical diversity of fungal secondary metabolites provides a significant advantage for the development of new drugs^{3,4}.

Anofinic acid, a potent active metabolite produced by *Aspergillus tubingensis*, has shown strong antimicrobial activity against a variety of hazardous bacteria in addition to its potent activity against various carcinoma cells, including Hep-G2, MCF-7, and HCT-116⁵. Sorbicillinoids possess the capability to be used as pharmaceutical and agrochemical agents due to their antiviral, antimicrobial, and anticancer effects besides their use as food

¹Department of Chemistry, Islamia College University, Peshawar, Khyber Pakhtunkhwa, Pakistan. ²Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, Khyber Pakhtunkhwa, Pakistan. ³Center of Chemical, Pharmaceutical and Food Sciences, Federal University of Pelotas, Pelotas RS, Brazil. ⁴Institute of Chemistry, Sao Paulo State University, Araraquara, Sao Paulo, Brazil. ✉email: drkhalidchem@yahoo.com; haroongold@gmail.com



Evaluation of potential bacterial protease inhibitor properties of selected hydroxyquinoline derivatives: an *in silico* docking and molecular dynamics simulation approach

Faiza Riaz^{a*}, Md. Sanower Hossain^{b*} , Miah Roney^{c*}, Yousaf Ali^d, Saira Qureshi^a, Riaz Muhammad^a, Said Moshawih^e , Shafida Abd Hamid^f, Veronique Seidel^g , Haroon ur Rashid^h and Long Chiau Ming^e

^aDepartment of Chemistry, Sarhad University of Science and Information Technology, Peshawar, Pakistan; ^bCentre for Sustainability of Ecosystem and Earth Resources (Pusat ALAM), Universiti Malaysia Pahang, Kuantan, Malaysia; ^cFaculty of Industrial Sciences and Technology, Universiti Malaysia Pahang, Kuantan, Pahang Darul Makmur, Malaysia; ^dFaculty of Allied Health Sciences, Iqra National University Swat Campus, Khyber Pakhtunkhwa, Pakistan; ^ePAP Rashidah Sa'adatun Bolkiah Institute of Health Sciences, Universiti Brunei Darussalam, Gadong, Brunei Darussalam; ^fKulliyyah of Science, International Islamic University Malaysia, Kuantan, Pahang, Malaysia; ^gNatural Products Research Laboratory, Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, UK; ^hInstitute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil

Communicated by Ramaswamy H. Sarma

ABSTRACT

Antimicrobial drug resistance (AMR) is a severe global threat to public health. The increasing emergence of drug-resistant bacteria requires the discovery of novel antibacterial agents. Quinoline derivatives have previously been reported to exhibit antimalarial, antiviral, antitumor, antiulcer, antioxidant and, most interestingly, antibacterial properties. In this study, we evaluated the binding affinity of three newly designed hydroxyquinolines derived from sulfanilamide (1), 4-amino benzoic acid (2) and sulfanilic acid (3) towards five bacterial protein targets (PDB ID: 1JIJ, 3VOB, 1ZI0, 6F86, 4CJN). The three derivatives were designed considering the amino acid residues identified at the active site of each protein involved in the binding of each co-crystallized ligand and drug-likeness properties. The ligands displayed binding energy values with the target proteins ranging from −2.17 to −8.45 kcal/mol. Compounds (1) and (3) showed the best binding scores towards 1ZI0/3VOB and 1JIJ/4CJN, respectively, which may serve as new antibiotic scaffolds. Our *in silico* results suggest that sulfanilamide (1) or sulfanilic acid (3) hydroxyquinoline derivatives have the potential to be developed as bacterial inhibitors, particularly MRSA inhibitors. But before that, it must go through the proper preclinical and clinical trials for further scientific validation. Further experimental studies are warranted to explore the antibacterial potential of these compounds through preclinical and clinical studies.

ARTICLE HISTORY

Received 13 August 2022
Accepted 6 November 2022

KEYWORDS

Antibacterial agents; hydroxyquinolines; molecular docking; ADMET; drug-likeness; *in silico*

Introduction

Infections remain one of the major threats to human health, and combatting infectious diseases has become a challenge worldwide (M. S. Hossain et al., 2021; S. Hossain et al., 2021; Sharma et al., 2018; Urbi et al., 2021). The human and economic costs of infections caused by bacteria, in particular, are a growing cause of concern (Muthukumar et al., 2019). Several bacterial pathogens have developed resistance to antibiotics, including strains identified as multi- or extensively-drug-resistant (El Faydy et al., 2017; Karuniawati et al., 2021). The Gram-positive bacterium Methicillin-resistant *Staphylococcus aureus* (MRSA) can cause skin, soft tissue, bone, joint infections, endocarditis and septicemia is a typical example of a drug-resistant microorganism (Qiu et al., 2001; Solomon & Oliver, 2014; Tjampakasari et al., 2021). Gram-negative bacteria also cause significant morbidity and

mortality worldwide; several strains have also become drug-resistant (Breijyeh et al., 2020).

Staphylococcus aureus infections are a global concern due to their high rate of developing antimicrobial resistance (AMR), limiting therapeutic options and healthcare-associated infections. The all-cause mortality rate, particularly from *S. aureus* bacteremia in the developed countries, reached up to 20–30%, with even over 50% rates in some less- and average-developing countries (Nakajima et al., 1978). The leading cause of AMR development is either one of them: (i) alteration of target, (ii) inactivation of antibiotic, (iii) bypass antibiotic and (iv) restricted antibiotic accumulation. Additionally, this process accelerated because of the misuse and overuse of antibiotics and poor infection prevention and control (Karuniawati et al., 2021; WHO, 2020). There are multiple enzymes involved in the development of AMR. For example,

CONTACT Md. Sanower Hossain  mshossainbge@gmail.com; mshossainbge@ump.edu.my  Centre for Sustainability of Ecosystem and Earth Resources (Pusat ALAM), Universiti Malaysia Pahang, Kuantan, 26300, Malaysia; Yousaf Ali  usaf_pk@yahoo.com  Faculty of Allied Health Sciences, Iqra National University Swat Campus, Khyber Pakhtunkhwa, Pakistan

*Faiza Riaz, Md. Sanower Hossain, and Miah Roney have contributed equally to this work, shared the first-author position, and ordered alphabetically.

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/07391102.2022.2146200>

RESEARCH

Open Access



Insights into metabolic and pharmacological profiling of *Aspergillus ficuum* through bioinformatics and experimental techniques

Zafar Ali Shah^{1,2}, Khalid Khan^{1*}, Haroon Ur Rashid³, Tanzeel Shah⁴, Mariusz Jaremko^{5*} and Zafar Iqbal²

Abstract

Background: Recently, numerous novel bioactive fungal metabolites have been identified that possess broad therapeutic activities including anti-inflammatory, antibiotic, antioxidant, and antitumor. The fungal mycochemicals as well as extracts have increased the interest of the scientific community in drug discovery research through a combination approach such as, molecular metabolic, pharmacological and computational techniques. Therefore, the natural fungus *Aspergillus ficuum* (*A. ficuum*) (FCBP-DNA-1266) was selected for metabolic and pharmacological profiling in this study.

Results: The metabolic profile of *A. ficuum* was explored for the first time and revealed the presence of bioactive compounds such as choline sulfate, noruron, hydroxyvittatine, aurasperone D, cetrimonium, kurilensoside, heneicosane, nonadecane and eicosane. Similarly, a pharmacological screen of *A. ficuum* was performed for the first time in *in vivo* and *in vitro* models. Interestingly, both the ethyl acetate and n-hexane fractions of *A. ficuum* were found to be more active against *Bacillus subtilis* among five tested bacteria with their zone of inhibition (ZOI) values of 21.00 mm $\pm$ 1.00 and 23.00 mm $\pm$ 1.00, at a concentration of 150 μ g mL⁻¹ respectively. Similarly, a significant decrease ($P < 0.001$) and ($P < 0.01$) in paw edema was observed in *A. ficuum*-treated animals at doses of 50 and 150 mg kg⁻¹, respectively, reflecting its potent anti-inflammatory effect. Furthermore, the docking results supported the antibacterial and anti-inflammatory effects of *A. ficuum*. In addition, the crude extract demonstrated no acute toxicity and the highest percent radical scavenging was recorded for both n-hexane and ethyl acetate extracts.

Conclusion: The metabolic profile of *A. ficuum* indicated the presence of biological relevant compounds. *A. ficuum* extract exhibited potent antibacterial and anti-inflammatory effects supported by docking results. Furthermore, *A. ficuum* extract demonstrated the highest percentage of radical scavenging activity along with no acute toxicity.

Keywords: Metabolic Profiling, Pharmacological screening, Molecular docking, *Aspergillus ficuum*

Introduction

Human life is under threat worldwide as mortality increases in developing countries. The spread and development of infectious diseases and drug-resistant pathogens has stimulated the search for new pharmaceutical agents [1]. The development of novel bioactive drugs is not only required against microbial resistance but also very important to overcome degenerative diseases. As a

*Correspondence: drkhalidchem@yahoo.com; Mariusz.jaremko@kaust.edu.sa

¹ Department of Chemistry, Islamia College, Peshawar, KP, Pakistan

⁵ Biological and Environmental Science and Engineering Division, King Abdullah University of Science and Technology, Thuwal, Makkah Province 23955-6900, Saudi Arabia

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Research



Cite this article: Iqbal A, Iqbal G, Umar MN, ur Rashid H, Khan SW. 2022 Synthesis of novel silica encapsulated spiropyran-based thermochromic materials. *R. Soc. Open Sci.* **9**: 211385. <https://doi.org/10.1098/rsos.211385>

Received: 26 August 2021

Accepted: 31 January 2022

Subject Category:

Chemistry

Subject Areas:

organic chemistry

Keywords:

thermochromism, spiro-heterocycles, spiropyran, colorimetric

Author for correspondence:

Anila Iqbal

e-mail: a.anilaiqbal@gmail.com

This article has been edited by the Royal Society of Chemistry, including the commissioning, peer review process and editorial aspects up to the point of acceptance.



Synthesis of novel silica encapsulated spiropyran-based thermochromic materials

Anila Iqbal¹, Ghazala Iqbal², Muhammad Naveed Umar³, Haroon ur Rashid⁴ and Sher Wali Khan⁵

¹Nanoscience and Technology Department, National Center for Physics, PO Box No 2141, Islamabad 44000, Pakistan

²Department of Pharmacy, Kohat University of Science and Technology, (KUST), Kohat, KP, Pakistan

³Department of Chemistry, University of Malakand, Chakdara Dir (L), KP, Pakistan


⁴Institute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil

⁵Department of Chemistry, Shaheed Benazir Bhutto University, Sheringal, Upper Dir, KP, Pakistan

A series of novel spiropyrans were synthesized through the condensation of substituted 3,3-dimethyl-2-methyleneindoline with different nitro-substituted *o*-hydroxy aromatic aldehydes. Indoles were initially substituted with a variety of alkanes and esters moieties. The substituted 3,3-dimethyl-2-methyleneindoline was then reacted with nitro-substituted *o*-hydroxy aromatic aldehydes to yield the respective spiropyrans. The synthesized novel spiropyrans were encapsulated in silica nano-shells to protect them from the effect of moisture and pH. The thermochromic behaviour of novel spiropyrans was studied by UV-visible spectroscopy. The thermally induced isomerization of spiropyran derivatives was carried out in a water/ethanol mixture. The thermal isomerization of spiro-heterocyclic (colourless form) to merocyanine (MC) (coloured form) was a discontinuous process and was observed in a temperature range of 5–60°C via UV-visible spectrometer. The absorption process occurs reversibly regardless of the heating/cooling sequence. The spiropyran derivatives, therefore, have a potential application for colorimetric temperature indication.

1. Introduction

The phenomenon of thermochromism has been known since 1926 when the reversible colour change was observed with temperature [1]. Therefore, thermochromism can be defined as the reversible thermo-transformation of chemical species between two forms



Molecular docking and dynamic simulations of Cefixime, Etoposide and Nebrodenside A against the pathogenic proteins of SARS-CoV-2

Haroon ur Rashid^{a,1,*}, Nasir Ahmad^{b,1}, Mohnad Abdalla^c, Khalid Khan^b,
Marco Antonio Utrera Martines^a, Samah Shabana^d

^a Institute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, 79074-460, Brazil

^b Department of Chemistry, Islamia College University, Peshawar, Khyber Pakhtunkhwa, Pakistan

^c Key Laboratory of Chemical Biology (Ministry of Education), Department of Pharmaceutics, School of Pharmaceutical Sciences, Cheeloo College of Medicine, Shandong University, 44 Cultural West Road, Shandong Province 250012, PR China

^d Department of Biochemistry and Molecular Biology, College of Marine Life Sciences, Ocean University of China, Qingdao 266003, PR China

ARTICLE INFO

Article history:

Received 17 May 2021

Revised 6 August 2021

Accepted 12 August 2021

Available online 13 August 2021

Keywords:

Coronavirus

Pathogenic protein

Etoposide

Cefixime

ABSTRACT

The catastrophe of the coronavirus continues from one part of the world to another, and hardly a country is left without its devastations. Millions of people were infected and several hundred thousand died of the COVID-19 pandemic across the world. There is no clear targeted drug therapy available for the treatment of the patients. The discovery of vaccines is not enough to curtail its spread and disastrous implications. An instantly qualifying approach is needed to utilize the current drugs and isolated compounds. The purpose of this work is to determine potent inhibitors against the target proteins of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). For this purpose, molecular docking study of pathogenic spike glycoproteins (S), nucleocapsid phosphoprotein (N), an envelope protein (E), two drugs i.e., cefixime, etoposide, and a previously isolated compound nebroadenside A is performed. Promising results were obtained via complimentary analysis of molecular dynamics (MD) simulations performed for the complexes of three proteins with etoposide drug. Minimum values were recorded for the docking scores and binding energies of the complexes. These results were further supported by the RMSD, RMSF data for the stability of proteins and ligands. Additionally, ligand properties and ligand-protein contacts were also explained with histograms of every simulation trajectory. The computational studies confirmed that cefixime, etoposide, and nebroadenside A can be used as potent inhibitors of COVID-19. Nevertheless, additional experimental investigations and validation of the selected candidates are mandatory to confirm their applicability for clinical trials.

© 2021 Elsevier B.V. All rights reserved.

Introduction

Coronavirus disease 2019 (COVID-19) is a contagious disease triggered by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). It was discovered in Wuhan, China, and has since outspread around the world like a pandemic, causing more than 150.6 million infections and about 32 hundred thousand deaths as of May 8, 2021 [1,2]. The World Health Organization called the disorder COVID-19 (coronavirus disease 2019). Genome sequencing reveals high similarity between SARS-CoV-2 and bat coronaviruses (CoV) especially beta-CoV and RaTG13 viruses that cause human diseases such as the Middle East respiratory syndrome (MERS) and

the Severe acute respiratory syndrome (SARS) and to a smaller extent HCoV-OC43 and HCoV-HKU1 [3,4]. The nucleocapsid protein, spike (S) protein, envelope (E) protein, membrane (M) protein, and coronavirus key protease participate actively in gene formulation and break polypeptides into replication-associated proteins. They are all encoded in the genome of the new coronavirus (SARS-CoV-2) [5]. The SARS-CoV-2 genome, like that of other coronaviruses, encrypt spike (S) glycoproteins, which project from the exterior of grown-up virions. The S glycoprotein is required for virus adhesion, incorporation, and infection of the host cell. The S glycoprotein's external spot makes it a straightforward target for the host immune feedbacks, making it a principal target for compensating the antibodies. The S protein is the subject of several vaccination campaigns and treatments due to its critical role in viral infection and innate immunity [6,7]. The nucleocapsid phosphoprotein (N) is a multifunctional and immunogenic virulence and pathogenicity determinant whose primary roles are oligomerization and packing

* Corresponding author.

E-mail addresses: haroongold@gmail.com (H.u. Rashid), mohnadabdalla200@gmail.com (M. Abdalla).

¹ These authors have contributed equally

Article

Conditions Optimization and Physiochemical Analysis of Oil Obtained by Catalytic Pyrolysis of Scrap Tube Rubber Using MgO as Catalyst

Riaz Muhammad ^{1,2,*} , Yousaf Ali ¹, Younes Messaddeq ³, Haroon ur Rashid ⁴ , Marco Antonio Utrera Martines ⁴ , Muhammad Naveed Umar ⁵, Sher Wali Khan ⁶ and Ali Riaz ^{1,3}

¹ Sarhad University of Science & Information Technology, Peshawar 25000, Pakistan; yousaf.chemistry@suit.edu.pk (Y.A.); ali.riaz.1@ulaval.ca (A.R.)

² Leibniz-Institute for Catalysis, University of Rostock, D-18059 Rostock, Germany

³ Centre for Optics, Photonics and Laser, University of Laval, 2375 rue de la Terrasse, Quebec, QC G1V 0A6, Canada; younes.messaddeq@copl.ulaval.ca

⁴ Institute of Chemistry, Federal University of Mato Grosso do Sul, Cidade Universitaria, Campo Grande 79074-460, MS, Brazil; haroongold@gmail.com (H.u.R.); marcomartines@gmail.com (M.A.U.M.)

⁵ Department of Chemistry, University of Malakand, Chakdara 18800, Pakistan; m.naveedumar@uom.edu.pk

⁶ Department of Chemistry, Shaheed BB University Sheringal, Sheringal 18050, Pakistan; sherwalikhn@yahoo.co.uk

* Correspondence: riazm.chemistry@suit.edu.pk; Tel.: +92-311-9830166



Citation: Muhammad, R.; Ali, Y.; Messaddeq, Y.; ur Rashid, H.; Antonio Utrera Martines, M.; Naveed Umar, M.; Khan, S.W.; Riaz, A. Conditions Optimization and Physiochemical Analysis of Oil Obtained by Catalytic Pyrolysis of Scrap Tube Rubber Using MgO as Catalyst. *Catalysts* **2021**, *11*, 357. <https://doi.org/10.3390/catal11030357>

Received: 27 January 2021

Accepted: 5 March 2021

Published: 10 March 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Motor vehicles scrap tires and tube rubbers generate a large amount of waste with different characteristics and compositions, contaminating the environment when not properly disposed. Waste inner tube rubber (isobutylene isoprene) representing a threat to the environment can be used as valuable source of energy. Waste inner tube rubber was pyrolyzed thermally under atmospheric pressure both with and without catalyst. Parameters of temperature, time, and catalyst weight were optimized for catalytic and thermal pyrolysis of isobutylene-isoprene rubber into liquid fuel, using MgO as catalyst. It was found that one-hour heating time at 350 °C using 2 g catalyst (MgO) constitutes a suitable parameter for the maximum conversion of scrap inner tube rubber into oil. The oil obtained was characterized by physical and chemical tests. Among the physical tests, Density, specific gravity, viscosity, kinematic viscosity, aniline point and flash point were determined according to IP and ASTM standard valves. The physical tests indicate the presence of aromatic and olefinic hydrocarbons. Among the chemical tests, the phenol test, bromine number, bromine water test, and KMnO₄ tests were conducted. The chemical tests are also the support of physical tests conducted. The physical and chemical tests indicate that the oil obtained is a mixture of kerosene, diesel, and light oil and could be used for fuel purposes.

Keywords: pyrolysis; scrap rubber; liquid fuel; magnesium oxide; catalysis

1. Introduction

Human lifestyle is improving rapidly, resulting in the increased demand for high-end freeways and highways, consequently increasing the demand for motor vehicles around the world. The rapid expansion of automobile business has therefore brought up some environmental drawbacks where handlings of waste tires/tubes are one of the fast-growing issues [1]. European Tyre & Rubber Manufacturers' Association (ETRMA) statistics reported that 1400 million tires are being produced, generating 17 million tons of used tires/tube waste every year [2].

Millions of tires/tubes are either thrown away or buried all over the world which is considered a threat to the ecosystem. As per a concise study, there would be 5000 million tires discarded annually by the year 2030 [3]. These scrap tire/tube rubber due to their

REVIEW

[View Article Online](#)
[View Journal](#) | [View Issue](#)Cite this: *RSC Adv.*, 2021, 11, 6060

Research developments in the syntheses, anti-inflammatory activities and structure–activity relationships of pyrimidines†

Haroon ur Rashid,^{ab} Marco Antonio Utrera Martinez,^{*a} Adriana Pereira Duarte,^{id a} Juliana Jorge,^a Shagufta Rasool,^b Riaz Muhammad,^b Nasir Ahmad^c and Muhammad Naveed Umar^{*d}

Pyrimidines are aromatic heterocyclic compounds that contain two nitrogen atoms at positions 1 and 3 of the six-membered ring. Numerous natural and synthetic pyrimidines are known to exist. They display a range of pharmacological effects including antioxidants, antibacterial, antiviral, antifungal, antituberculosis, and anti-inflammatory. This review sums up recent developments in the synthesis, anti-inflammatory effects, and structure–activity relationships (SARs) of pyrimidine derivatives. Numerous methods for the synthesis of pyrimidines are described. Anti-inflammatory effects of pyrimidines are attributed to their inhibitory response *versus* the expression and activities of certain vital inflammatory mediators namely prostaglandin E₂, inducible nitric oxide synthase, tumor necrosis factor- α , nuclear factor κ B, leukotrienes, and some interleukins. Literature studies reveal that a large number of pyrimidines exhibit potent anti-inflammatory effects. SARs of numerous pyrimidines have been discussed in detail. Several possible research guidelines and suggestions for the development of new pyrimidines as anti-inflammatory agents are also given. Detailed SAR analysis and prospects together provide clues for the synthesis of novel pyrimidine analogs possessing enhanced anti-inflammatory activities with minimum toxicity.

Received 19th December 2020
Accepted 27th January 2021

DOI: 10.1039/d0ra10657g

rsc.li/rsc-advances

1. Introduction

The word inflammation is derived from the Latin “flamma” meaning flame. It is normal feedback of the body to safeguard tissues against disease or infection. The inflammatory reaction initiates with the generation and discharge of chemical agents from the cells in the diseased, infected, or wounded tissue. Inflamed tissues produce extra signals that use leukocytes (white blood cells) at the position of inflammation. Leukocytes damage any infective or harmful agent and eliminate cellular residues from damaged tissue. This inflammatory reaction generally stimulates the curing process. However, an uncontrolled inflammatory response may become detrimental.^{1,2} Inflammation represents a portion of the body's immune response. An inflammatory reaction is liable for curing wounds,

infections, and any damage to the tissues. Numerous feedbacks produced by the defense mechanism as a response to a physical injury or an infection result in inflammation.³ Acute inflammation grows rapidly and turns out to be severe in a brief time. Its symptoms continue for a few days but may last for several weeks under certain circumstances. Common indications of acute inflammation include swelling, redness, pain, immovability, and heat. Acute inflammation can be caused by certain conditions and diseases including acute bronchitis, abrasion or cut on the skin, achy throat from flu or cold, infected ingrown toenail, acute appendicitis, sinusitis, tonsillitis, dermatitis, infective meningitis, high-intensity exercise, and physical trauma.^{4,5} Chronic inflammation persists for a prolonged duration (for several months and even years) in which active inflammation, tissue damage, and repair occur concurrently. Usually, the extent and effects of chronic inflammation differ with the source of the injury and the efficiency of the body to heal and control the damage. Chronic inflammation is characterized by typical symptoms such as body pain, fever, rash, weight gain or weight loss, fatigue, joint pain, and mouth sores. Chronic inflammation may lead to the progression of certain diseases namely diabetes, cancer, cardiovascular diseases, rheumatoid arthritis, allergies, chronic obstructive pulmonary disease (COPD), tuberculosis, asthma, hepatitis, periodontitis, and chronic peptic ulcer.^{1,6} Fig. 1 indicates the health

^aInstitute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil. E-mail: marcomartines@gmail.com

^bDepartment of Chemistry, Sarhad University of Science and Information Technology, Peshawar, Khyber Pakhtunkhwa, Pakistan

^cDepartment of Chemistry, Islamia College University, Peshawar, Khyber Pakhtunkhwa, Pakistan

^dDepartment of Chemistry, University of Malakand, Chakdara, Dir (L), Khyber Pakhtunkhwa, Pakistan. E-mail: m.naveedumar@uom.edu.pk

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ra10657g

RESEARCH ARTICLE

Discovery of Biomarkers in Hepatocellular Carcinoma Metastasis Using Bioinformatic Analysis

Jinrui Wei^{1,2}, Haroon ur Rashid³, and Lichuan Wu^{1,*}

¹Medical College of Guangxi University, Nanning, 530004, China; ²Guangxi Scientific Research Center of Traditional Chinese Medicine, Guangxi University of Chinese Medicine, Nanning, Guangxi, China; ³Institute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil

Abstract: Background: Liver cancer is one of the most deadly malignancies worldwide. Tumor metastasis is the main cause of liver cancer-related death. So far, the mechanism of liver cancer metastasis is far away from fully elucidated. In this study, we aimed to discover key regulators involved in liver cancer metastasis by data mining.

Methods: Two different types of data, including mRNA microarray (GSE6222 and GSE6764) and miRNA microarray (GSE67138), were analyzed. A total of 83 intersectant differently expressed genes (DEGs) with the same expression pattern in GSE6222 and GSE6764 were identified. One hundred and thirty-one differently expressed miRNAs (DEMs) were identified in GSE 67138. Furthermore, a total of 26 pairs of miRNA-target, including 18 DEMs and 13 DEGs were identified as critical miRNA-target axis via miRNA-target gene interaction analysis.

Results and Conclusion: Among the 18 DEMs and 13 DEGs, 10 miRNAs and 10 target genes are significantly correlated with patients' survival ($p < 0.05$). Our results and methods might be interesting for data mining and helpful for further experimental functional validation.

ARTICLE HISTORY

Received: Feb 09, 2020

Revised: April 10, 2020

Accepted: May 07, 2020

DOI:

10.2174/1574893615999200713163643

Keywords: Bioinformatic analysis, biomarkers, hepatocellular carcinoma, metastasis, liver cancer, key genes.

1. INTRODUCTION

Liver cancer is one of the most prevalent malignancies, which is the fourth common cause of mortality worldwide [1]. Liver cancer presents a heavy burden for many countries due to its high aggressiveness and poor survival. Tumor metastasis is the leading cause of liver cancer-related death [2]. Presently, the mechanism of HCC metastasis has not been fully elucidated. Thus, in-depth understanding of HCC metastasis can be useful in clinical settings for prognosis and treatment.

Aberrant expressions of gene and dysregulated signaling pathways are key hallmarks of cancer. It has been reported that cancer cells may benefit from undesirable changes in microRNAs (miRNAs) [3]. It is well-recognized that miRNAs play a vital role in the cancer metastasis by negatively regulating their target genes [4]. Han reported that miR-532-3p could regulate KIFC1 to promote metastasis in HCC [5]. Xu reported miR-876-5p could inhibit HCC metastasis by targeting BCORL1 [6]. These results implied that the

characterization of dysregulated miRNAs and their target genes could uncover potential metastasis biomarkers of HCC.

In recent years, with the rapid development in high-throughput technologies, gene microarray appears to be an effective method for quickly detecting differentially expressed genes (DEGs) between tumor and normal samples [7]. A large number of data have been generated and stored in public platforms, such as the Gene Expression Omnibus (GEO) profiles, which could be accessed for free [8]. Although prognostic biomarkers of HCC have been identified by analyzing datasets from GEO [9], results are always inconsistent due to sample heterogeneity, leading to unreliable results. To get reliable results, an integrated bioinformatics analysis of different types of data could be applied. In the present study, two gene microarray and one miRNA microarray datasets were retrieved and analyzed with R software (version 3.4.4) and Bioconductor packages (<http://www.bioconductor.org/>) to produce DEGs and DEMs. Then, miRNet and Kaplan-Meier Plotter online databases were used to construct miRNA-protein interactions and predict potential miRNAs-targets pairs in HCC (Fig. 1). Our results and method might be interesting for data mining and helpful for further experimental functional validation.

*Address correspondence to this author at the Medical College of Guangxi University, Nanning, 530004, China; Guangxi Scientific Research Center of Traditional Chinese Medicine, Guangxi University of Chinese Medicine, Nanning, Guangxi, China;
E-mail: richard_wu@gxu.edu.cn, lichuanwu@gmail.com


Review | <http://dx.doi.org/10.17807/orbital.v13i4.1638>

Potential Drug Candidates in Clinical Trials for the Treatment of Covid-19: An Updated Overview

Osmar Ignacio Ayala Cáceres [✉] †, Fernanda Timóteo [✉] †, Kristiane Fanti Del Pino Santos [✉], Rafael Rodrigo Piva Vasconcelos [✉], Marco Antonio Utrera Martines [✉], Juliana Jorge [✉], and Haroon ur Rashid [✉]

The coronavirus (SARS-CoV-2) emerged in December 2019 in Wuhan, China and by the end of April 2020, it spread to more than 200 countries worldwide. Particularly the USA, Brazil, Spain, Italy, France, Germany, UK, Turkey, Iran, and Russia were the most affected countries till December 2020. Currently, most of the researchers are in a continuous struggle to develop a vaccine or new drugs to combat Covid-19. There are more than 30 drug candidates including Western medicines, natural products, and traditional Chinese medicines, that have shown to exhibit some efficacy against this highly infectious virus. This review encompasses the potential efficacy of some key drugs recently tested against Covid-19. With the rapid spread of Covid-19 reaching a new level every day, there is an immediate need to find safe and effective measures to diagnose, treat, mitigate, and combat the disease. Looking at the alarming dimensions that the disease is acquiring, treatment strategies among the different drug systems are being investigated. Currently, clinical management includes infection prevention, control measures, and supportive care, including supplemental oxygen and mechanical ventilation when indicated. The pharmaceutical interventions evaluated for the treatment of Covid-19 include human immunoglobulin, interferons, chloroquine, hydroxychloroquine, arbidol, remdesivir, oseltamivir, favipiravir, carrimycin, methylprednisolone, bevacizumab, thalidomide, vitamin C, pyrofenidone, darthviravir, bromexistone, ryrexavir, lopinavir, xianping, and traditional Chinese medicine. But still, the researchers are struggling to discover the most effective drug for the treatment of the current pandemic of Covid-19.

Graphical abstract



Keywords

Coronavirus
Genome
Nucleocapsid
Pandemic
Protease inhibitor

Article history

Received 04 July 2021
Revised 05 August 2021
Accepted 15 August 2021
Available online 30 September 2021
month year

Handling Editor: Adilson Beatriz

Institute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil. †These authors have contributed equally. Corresponding author. E-mail: ju.jorge@yahoo.com.br (J. Jorge); haroongold@gmail.com (H. ur Rashid).

EMODIN-CONTAINING MCM-41 TYPE MESOPOROUS SILICA NANOPARTICLE DRUG DELIVERY SYSTEM

I. M. D. dos Santos¹, S. P. Pereira¹, A. V. Mezacasa¹, O. I. A. Caceres¹,
F. Timoteo¹, A. S. Lopes¹, K. F. del Pino¹, A. P. Duarte¹,
T. F. M. Cardoso², G. R. de Castro³,
H. U. Rashid¹, and M. A. U. Martinez¹

UDC 544.23.022

With the aim of developing drug delivery systems, it was shown that nanoparticles of mesoporous silica molecular sieve MCM-41 with average size 100-150 nm and specific surface 655 m²/g could include 202 mg of emodin per one gram of the carrier and release more than 85% of this compound at pH 8 during 48 h.

Keywords: Emodin, anthraquinone, mesoporous silica nanoparticles, drug controlled release system, ultraviolet-visible

Emodin (Fig. 1), also known as 1,3,8-trihydroxy-6-methyl-anthraquinone, a natural compound extracted from herbaceous plants such as *Reynoutria japonica* and *Rheum officinale*, has been widely used as a traditional medicine in some regions of Asia [1]. It has been reported to possess biological activities such as antibacterial, anti-inflammatory, immunosuppressive, vasodilator, laxative, and antineoplastic [2, 3].

One major drawback of emodin as a biological agent is its low solubility in water [1]. The biological system consists of different types of barriers with hydrophobic or hydrophilic characteristics. The biodistribution of a drug in an organism is related to its capability to pass through tissues containing aqueous and lipidic environments. The majority of drugs are organic and generally have poor water solubility, which in turn interferes with their biodistribution and affects the efficiency of cell delivery [4].

In this perspective, substantial efforts have been made to discover more efficient and suitable liberation system, so that drugs can reach their targets in the estimated time and thus can ensure the desired effects only. Therefore, in pursuit of some potential strategies to promote a long-lasting release of insoluble-water drugs, emodin has been used as a model drug for research on controlled liberation systems (CLS) [5, 6].

The research field of CLS is in focus among the scientific community and is proliferating, with nanotechnology playing a pivotal role in this expansion [7].

¹Institute of Chemistry, Federal University of Mato Grosso do Sul, Cidade Universitaria, 79074-460 Campo Grande, MS, Brazil. E-mail: marco.martines@ufms.br.

²Faculty of Pharmaceutical Sciences, Food and Nutrition, Federal University of Mato Grosso do Sul, Cidade Universitaria, 79070-900 Campo Grande, MS, Brazil

³Institute of Biosciences, Sao Paulo State University, 18618-689 Botucatu, SP, Brazil



Anti-cancer potential of sophoridine and its derivatives: Recent progress and future perspectives

Haroon ur Rashid^{a,b,*}, Shagufta Rasool^b, Yousaf Ali^b, Kamin Khan^b,
Marco Antonio Utrera Martinez^{a,*}

^a Institute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil

^b Department of Chemistry, Sarhad University of Science and Information Technology, Peshawar, Khyber Pakhtunkhwa, Pakistan

ARTICLE INFO

Keywords:

Sophoridine
Phytochemical
Alkaloid
Cancer
Topoisomerase I

ABSTRACT

Cancer is the second leading cause of mortality and has resulted in about 9.6 million deaths around the world in 2018. Cancer-caused deaths are expected to be 11.5 million by 2030 all over the world. Because of the fatal nature of cancer, substantial efforts are made all over the world to combat it. Phytoconstituents such as certain alkaloids, saponins, tannins, polyphenols, and terpenoids exhibit anticancer effects. Sophoridine is a tetracyclic quinolizidine alkaloid isolated from the stem and leaves of medicinal plants *Sophora alopecuroides* L., and *Euchresta japonica* Benth, and roots of *Sophora alopecuroides* Ait. Chinese Food and Drug Administration (CFDA) approved sophoridine as an antitumor agent in 2005. This review covers the antitumor activities of sophoridine and its derivatives. The efficacy of sophoridine analogs is expressed with respect to their half-maximal inhibitory concentration (IC₅₀ values). Structure-activity relationship (SAR) study for most of the sophoridine derivatives has been explained. Moreover, the current market of anticancer drugs and its expected growth are discussed. Prospects provide suggestions and clues for novel sophoridine-based anticancer agents with enhanced expected efficacy and minimum toxicity.

1. Introduction

Cancer represents the uncontrolled division of abnormal body cells. Cancer cells attack or reach to the adjacent healthy tissues of the body via the bloodstream and lymphatic systems, causing malignant tumors. On the contrary, cells in benign tumors neither invade the neighboring tissues nor spread to other body parts. Nearly 90% of cancer-related mortalities are triggered by tumor dispersion, a phenomenon termed as metastasis. (Fig. 1) [1,2] DNA damage is reckoned to be the primary cause of cancer growth. It produces mutations and epimutations, which then lead to cancer progression through the process of natural selection. (Fig. 2) [3] Cancer is the second-highest cause of deaths in the world, only excelled by cardiovascular ailments [4].

According to a recent press release of the International Agency for Research on Cancer (IARC), 18.1 million new cancer cases, and 9.6 million cancer-caused deaths were reported worldwide in 2018 [5]. According to a report from the World Health Organization (WHO), over 70% of the cancer-related mortalities happened in developing countries. Cancer-related deaths will increase and are expected to reach an approximate number of 11.5 million by 2030 all over the world. Major

risk factors for cancer include alcohol consumption, smoking, dietary aspects (comprising inadequate vegetables and fruits consumption), obesity, lack of physical activity, exposure to ionizing and non-ionizing radiations, and exposure to toxins, chronic infections from helicobacter pylori, human papillomavirus (HPV), hepatitis B virus (HBV), and hepatitis C virus (HCV) [6].

Approximately 100 kinds of cancer have been recognized, which are characteristically named for the organs or tissues where they arise. Statistics show that prostate and breast cancers are the two principal kinds of cancer identified in males and females, respectively [1,7]. In modern medicine, surgery, radiotherapy, and systemic therapy (such as chemotherapy, immunotherapy, hormonal therapy, and targeted therapy) represent the key modalities applied to treat different types of cancer [8,9]. Owing to the deadly nature of cancer, considerable efforts are made worldwide to combat it by the exploration of novel antitumor agents, which has resulted in a noteworthy success. US Food and Drug Administration (FDA) recommended approximately 150 antitumor drugs from 1949 to 2014. Depending on their mode of action, these drugs have been classified into two sets (a) 61 cytotoxic drugs and (b) 89 target-oriented drugs [10–13]. Selected FDA approved drugs are

* Corresponding authors at: Institute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil (H. ur Rashid).

E-mail addresses: haroongold@gmail.com (H. ur Rashid), marcomartines@gmail.com (M.A.U. Martinez).



New modification strategy of matrine as Hsp90 inhibitors based on its specific L conformation for cancer treatment

Yiming Xu^{a,c,1}, Dewang Jing^{c,1}, Dong Zhao^b, Yongji Wu^d, Lu Xing^b, Haroon ur Rashid^{c,e}, Haodong Wang^c, Lisheng Wang^{a,*}, Huiling Cao^{b,*}

^a Medicinal College, Guangxi University, Nanning 530004, China

^b Shaanxi Key Laboratory of Ischemic Cardiovascular Disease, Institute of Basic Translational Medicine, Xi'an Medical University, Xi'an 710021, China

^c School of Chemistry and Chemical Engineering, Guangxi University, Nanning 530004, China

^d College of Veterinary Medicine, Northwest A&F University, Yangling 712100, China

^e Department of Chemistry, Sarhad University of Science & Information Technology, Peshawar, Khyber Pakhtunkhwa 25120, Pakistan

ARTICLE INFO

Keywords:

L-shaped matrine derivatives
Radicicol
Thermal shift assay
Anticancer activity
Mechanism of action

ABSTRACT

The similarity of spatial structure between radicicol and matrine urged us to perform conformation modification of matrine, followed by L-shaped matrine derivatives, **6**, **12**, **21a-h** and **22a-h** were originally designed, synthesized and evaluated for Hsp90^N inhibitors as anticancer agents. TSA (Thermal Shift Assay) results indicated that **21e**, **22a-c** and **22e-g** exhibited strong binding force against Hsp90^N with $|\Delta T_m| > 3$, meanwhile, MTT assay also revealed these compounds displayed potent anticancer activity with IC₅₀ values below 25 μ M against HepG2, HeLa and MDA-MB-231 cells lines. Then, compound **22g** with a high $\Delta T_m = 10.92$ was chosen as a representative to perform further mechanism study. It can induce cell apoptosis, arrest the cell cycle at the S phase and decrease the expression level of Hsp90 in Hela cell. These results originally provided targeted modification strategy for matrine derivatives to serve as Hsp90 inhibitors for cancer therapy.

1. Introduction

Matrine (Fig. 2d), a quinolizidine natural product extracted from *Sophora flavescens* (Kushen),^{1–4} was approved by Chinese FDA in 1995 for cancer treatment. Currently, matrine has been widely used to adjuvant treatment of liver cancer, lung carcinoma and gastric cancer with low side effects for many years in China.^{5,6} Although large-scale application of matrine with reference to its use for the treatment of cancer was confirmed,^{7–11} its activity is moderate, which might be caused by its speculated features such as broad binding sites but low specificity. Thus, enhancing the sensibility and specificity against anticancer targets of matrine attracted our attention. On the other hand, matrine possess many advantages such as high solubility, good safety profiles and a special chemical scaffold, suggesting that it is an ideal lead compound for further modifications and optimizations to treat cancer.

Recently, our group also conducted a comprehensive literature review (Fig. 1) on modification strategy of matrine.¹² As illustrated in Fig. 1, lots of matrine derivatives have been synthesized through various modifications on different positions of matrine. However, up to now, research on targeted design for matrine derivatives is still not be involved

which caused blindness for their development. Different from traditional cytotoxic chemotherapy, targeted therapy has many advantages, such as high specificity, good efficacy and low toxicity. Inspired by above suggestions, targeted design for matrine might be an effective strategy to discover anticancer candidates of matrine derivatives.

Interestingly, the specific L-shaped conformation of matrine is similar with radicicol (Fig. 2). Besides, radicicol serves as a Hsp90^N inhibitor and its molecular mechanism has been thoroughly studied, which provided possibility for structural modification of matrine derivatives to target Hsp90^N based on radicicol conformation analysis. Moreover, the overexpressed characteristics in tumor cells and the prospect of stalling a number of oncogenes by inhibiting a single protein have made Hsp90 an attractive target for cancer therapy.^{13–15} Thus, by targeted design and inhibiting the activity of Hsp90, it might be an effective strategy for matrine modifications to treat cancer.

2. Drug design

Radicicol, as the Hsp90^N inhibitor, its mechanism involves a competition with ATP at N-terminal ATP-binding pocket of Hsp90, resulting

* Corresponding authors.

E-mail addresses: lswang@gxu.edu.cn (L. Wang), caohuiling_jzs@xjtu.edu.cn (H. Cao).

¹ These authors contributed equally to this work.

SYNTHESIS AND CHARACTERIZATION OF GADOLINIUM-BASED NANOCRYSTALS FOR THEIR POTENTIAL APPLICATION AS TRIMODAL IMAGING CONTRAST AGENTS

HAROON UR RASHID^a, **MUHAMMAD YASEEN**^b, **ALI RIAZ**^{a*}, **KAMIN KHAN**^a, **HANIF UR RAHMAN**^a,
KHALID KHAN^c, **NASIR AHMAD**^c AND **MARCO ANTONIO UTRERA MARTINES**^{d*}

^aDepartment of Chemistry, Sarhad University of Science & Information Technology, Peshawar, Khyber Pakhtunkhwa, Pakistan.

^bInstitute of Chemical Sciences, University of Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan.

^cDepartment of Chemistry, Islamia College University, Peshawar, Khyber Pakhtunkhwa, Pakistan.

^dInstitute of Chemistry, Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil.

ABSTRACT

Lanthanide-doped gadolinium nanomaterials have attracted considerable attention due to their promising applications in biological imaging. Sodium gadolinium fluoride (β -NaGdF₄) nanomaterials act as perfect host for doping of luminescent lanthanide ions. Due to heavy nature of gadolinium, it can strongly absorb X-ray radiations. Therefore, Gd-based nanomaterials are applied as contrast agents for X-ray tomography (CT). Presence of seven unpaired electrons in Gd³⁺ ion, its large magnetic moment and long electronic relaxation time makes it an ideal candidate to enhance water proton relaxation rates. These characteristics make Gd-based nanomaterial useful for their potential application as MRI contrast agents. In this work, sodium gadolinium fluoride is doped with ytterbium and erbium to produce β -NaGdF₄:Yb/Er as core nanocrystals. They are subsequently coated with sodium gadolinium fluoride doped with neodymium (β -NaGdF₄:Nd) to produce β -NaGdF₄:Yb/Er@ β -NaGdF₄:Nd core-shell nanocrystals. Powder X-ray diffraction (XRD) results indicated that β -NaGdF₄ (host), β -NaGdF₄:Yb/Er (core) and β -NaGdF₄:Yb/Er@ β -NaGdF₄:Nd (core-shell) nanocrystals all exist in hexagonal crystalline phase. While Dynamic Light Scattering (DLS) results indicated that the particle size of the synthesized nanocrystals was in range of 12 to 27 nm. FTIR results confirmed that the synthesized nanocrystals were stabilized by oleic acid. Nanocrystals reported in this work are expected to be useful trimodal contrast agents for photoluminescence, X-ray tomography and Magnetic Resonance Imaging (MRI).

Keywords: lanthanide, nanocrystals, trimodal, gadolinium, contrast agents.

INTRODUCTION

Recently, rapid research developments in the field of medicine and life science have also driven the biomedical research. This emerging and growing research field comprises of a number of bioassay approaches such as X-ray Computed Tomography (also called Computed Tomography, CT), Single-Photon Emission Computed Tomography (SPECT) [1-4], Positron Emission Tomography (PET) [5, 6] and Magnetic Resonance Imaging (MRI) [7-9]. These tomographic imaging methods have high penetration power. Therefore, they play a key role in biomedical imaging. However, these techniques have some unavoidable drawbacks such as limited spatial resolution, long scan acquisition time and application of strong ionizing radiations (except for MRI) [10-13]. Furthermore, these modalities differ from each other in spatial and temporal resolution, imaging depths, anatomical and molecular details and the necessary material properties of contrast agents essential to improve the quality of images in terms of their visibility. In many cases, it is beneficial to use several complimentary imaging techniques for rapid and precise diagnosis. Since majority of these imaging modalities apply exogenous contrast agents to improve the signal-to-noise ratio, therefore, the development of multimodal contrast agents is highly desirable to obtain images of better quality. Such agents have proved more useful for patient care.

Moreover, they also reduce the cost and increase the safety by restricting the number of contrast agents administration needed for imaging purposes [14]. The idea of designing multimodal contrast agents provides the prospective to integrate the benefits of different modalities while at the same time overcoming the shortcomings of one another [15, 16]. In addition, multimodal contrast agents can be very useful for the technical development of emerging imaging techniques [17-19]. Current research indicate that nanomaterial-based multimodal imaging probes have been developed for accurate imaging and diagnosis while applying different imaging modalities. Recently, lanthanide metal ions doped with upconverting (Excitation at lower energy than emission) nanomaterial have been investigated for their potential applications in biomedical imaging. Such nanomaterials possess high photostability and show neither photobleaching nor photoblinking [20]. Additionally, upconverting nanomaterials are excited by near-infrared (NIR) radiations which ensure optical imaging of deeper tissues

with significantly controlled autofluorescence, and create enhanced image contrast [21-23]. β -NaGdF₄ host nanomaterial doped with lanthanide ions combination of Yb³⁺, Er³⁺ or Yb³⁺, Tm³⁺ have exhibited significant upconversion luminescence [24]. Gadolinium is a heavy metal of high atomic number which strongly absorbs X-rays. Its trivalent ion (Gd³⁺) causes enhancement of water proton relaxation rate due to maximum number (seven) of unpaired electrons (among all lanthanide ions), a long electronic relaxation time ($\sim 10^{-9}$ s) and a large magnetic moment (7.9 BM). These properties make Gadolinium the best candidate for applications as luminescence, CT and MRI contrast agents [11, 25, 26]. Upconverting nanomaterial based multimodal contrast agents can serve for fluorescence imaging, CT and MRI modalities. Such contrast agents possess the advantages of high sensitivity (fluorescence imaging) as well as the potential of high spatial resolution imaging (of MRI and CT techniques) [27]. This work reports the synthesis of β -NaGdF₄ nanocrystals doped with Yb³⁺ and Er³⁺ (NaGdF₄:Yb/Er). They were synthesized as core nanocrystals and were subsequently coated with a shell of NaGdF₄:Nd³⁺ (Core-shell nanocrystals).

EXPERIMENTAL

Chemicals: Gadolinium Oxide (99.9%), Ytterbium(III) acetate hydrate (99.99%), Erbium Oxide (99.9%), Neodymium(III) acetate hydrate (99.9%), Sodium hydroxide ($\geq 97\%$), 1-octadecene (90%), oleic acid (90%), Ammonium fluoride ($\geq 99.9\%$), Acetic Acid glacial ($\geq 99.85\%$) and cyclohexane were purchased from Sigma-Aldrich. All chemicals were used without any further purification.

Synthesis of Gadolinium acetate hydrate

Gadolinium acetate hydrate was prepared following the procedure reported by Dühnen and Haase for the synthesis of gadolinium trifluoroacetate (Gd-TFA) [28]. Gd₂O₃ (3.63 g, 10 mmol) was added to a solution of 80 mmol of concentrated acetic acid (4.6 mL) and 12.4 mL of distilled water. The solution was heated under reflux (at ca. 90 °C) for 60 min to dissolve the Gd₂O₃ completely. After removing excess acetic acid and water with a rotary evaporator at 50 °C, Gadolinium acetate hydrate was obtained as a white powder. Subsequently the powder was dried at 60 °C for 17 h.